

## A Review of Subject

*Dr Nishith Govil<sup>1</sup>, Dr Priyanka Mishra<sup>2</sup>, Dr Amanta Lucy Ittoop<sup>3</sup>, Dr Prakash Chandra<sup>4</sup>*

1. *Dr Nishith Govil MBBS MD Department of Anaesthesiology AII INDIA INSTITUTE OF MEDICAL SCIENCES Rishikesh*
2. *Dr Priyanka Mishra MBBS MD Department of Anaesthesiology AII INDIA INSTITUTE OF MEDICAL SCIENCES Rishikesh*
3. *Dr Amanta Lucy Ittoop MBBS Department of Anaesthesiology AII INDIA INSTITUTE OF MEDICAL SCIENCES Rishikesh*
4. *Dr Prakash Chandra MBBS Department of Anaesthesiology AII INDIA INSTITUTE OF MEDICAL SCIENCES Rishikesh*

**Address for Correspondence: [editojohp@gmail.com](mailto:editojohp@gmail.com)**

**INTRODUCTION**

Hyperparathyroidism may be secondary to parathyroid hyperplasia, adenoma or carcinoma. Increased secretion of parathyroid hormone results in increased bone resorption, increased intestinal absorption of calcium, and decreased renal clearance of calcium. This will lead to increased serum calcium concentration which is responsible for most of the manifestations of the disease.

***Access this Article Online***

Website: <http://www.journalofhospitalpharmacy.in> Quick Response Code:

Received on 30/10/2018

Accepted on 12/11/2018 © HEB All rights reserved

